

# Ruffed Grouse Drumming Survey

## 2008

By Brian Dhuey

### **Abstract**

Statewide ruffed grouse population indices increased 7% between 2007 and 2008, based on the number of drumming grouse heard during roadside surveys. Changes in indices to breeding grouse populations varied by region, but the statewide mean number of drums/stop did not change significantly ( $P= 0.50$ ) from 2007 to 2008. Drummer densities on the Sandhill Wildlife Area in Wood County increased 8%. Stone Lake Experimental Area in Oneida County increased 4% from 2007 levels.

### **Methods**

#### **Statewide**

Counts of drumming grouse heard along roadsides were conducted on 89 transects throughout the state in 2008. Twenty-six statewide transects were considered to be zero and not run in 2008. This roadside survey has been conducted annually since 1964 by DNR wildlife managers, wildlife technicians, foresters, law enforcement personnel, USFS staff, and Ruffed Grouse Society volunteers to determine grouse population trends throughout Wisconsin. A new 10-stop survey on 117 randomly located transects was initiated in 1994 and continued in 2008. This year marked the fifteenth year that the "new" ruffed grouse surveys were run. As per the change over plan, no "old" drumming routes were run since 1996. Also, "new" routes which had counts of zero for the first three years were not to be run for three years. After that three year period, they are to be run again every 3 years to confirm that they indeed are still zero.

Procedures for the "new" routes were similar to the earlier survey protocols except for one count instead of two and 10 stops instead of 15. Survey data were entered into the DNR UNIX computer system and summarized using the Statistical Analysis System (SAS).

#### **Research Census Areas**

DNR research personnel have conducted a census of drummers on Sandhill Wildlife Area and Stone Lake Area since 1968. This survey has provided comparative statistics on population trends and an estimate of drummer density. Searches for males were conducted during favorable weather between 1 April and 10 May. The census on the Sandhill Wildlife Area encompassed 2,020 acres of grouse habitat in the area open to hunting and 1,300 acres within the un hunted portion of the area. The census on the Stone Lake Experimental Area in Oneida County encompassed 3,310 acres of grouse habitat.

## Results

### Statewide

Responses were received from wildlife managers, wildlife technicians, and other cooperators that helped conduct the survey on 89 survey transects in 2008. Twenty-six transects were considered to be zero and not run in 2008, and 2 transects that were not surveyed. The total number of routes used in estimating a statewide ruffed grouse drumming index in 2008 was 115. This is one less than the total number of transects (116) used in the spring of 2007.

Statewide, ruffed grouse population indices increased in Wisconsin between 2007 and 2008 (Table 1). This is the third increase in the ruffed grouse index in the last 3 years. Survey indices show an increase in drumming grouse in two of the four regions of the state (Fig. 1). Statewide, overall density increases were not significant ( $P= 0.50$ ) during 2007-2008. Transects completed in both 2007 and 2008 were compared to detect population changes. Transects were considered to have changed from last year if the change was greater than two drums per transect. The number of transects with increased drumming outnumbered by 33 to 23 those that showed decreases, with 59 transects unchanged.

Brood production in the summer of 2007 was slightly lower than in 2006 and may have set the stage for the decrease in breeding grouse in 2008. Wisconsin's primary grouse range, the Northern region, showed an increase in breeding grouse, it would appear that Wisconsin is still on the upswing of the current grouse cycle.

Survey conditions for 2008 were better than those in 2007. Surveyors rated the overall survey conditions as "excellent" on 60% of the transects run, while 49% rated the overall conditions as "excellent" in 2007. Surveyors rated the conditions as "Fair", the lowest available weather condition rating, 3% of the time in 2008 compared to 6% in 2007. The percent of observers rating the conditions as "excellent" was still below the highest level of 68% recorded in 1998. Survey conditions do influence drumming activity and may cause grouse numbers to be over or under estimated.

### Research Census Areas


Grouse numbers on the Sandhill Wildlife Area were up in 2008 (Table 2). Sandhill Wildlife Area increased 8% (43 vs. 40 birds in 2007) while the central region of the state showed a decrease of 3% in drumming activity. The unhunted portion of the wildlife area (1,300 acres) increased by 2 birds in 2008 (19 vs. 17 in 2007). The hunted portion of the wildlife area (2,020 acres) had a slight increase in breeding grouse, with 24 birds counted in 2008, up from 23 in 2007. The Stone Lake census area was again surveyed in 2008, seventy-four birds were counted this spring, seventy-one birds were counted in the spring of 2007. This is an increase of 4%, the northern region of the state showed a 12% increase on the road side survey. The survey technique used to measure grouse densities on these two areas is different than that used on the statewide survey. Any comparison of these results to statewide totals should be done cautiously.

**Table 1.** Ruffed Grouse drumming results 2007-2008, drums per stop (routes run), % change, and number of routes with a change of greater than 2 drums per route from 2007 levels.


Region	Drums/Stop 2007 (routes run)	Drums/Stop 2008 (routes run)	% Change	# of Decreasing Routes	# of Increasing Routes	# of Routes with No Change
Central	0.95 (27)	0.92 (27)	-3%	6	7	14
Northern	1.48 (42)	1.66 (42)	+12%	11	22	9
Southeast	0.09 (30)	0.12 (30)	+33%	4	2	24
Southwest	0.46 (17)	0.37 (16)	-20%	2	2	12
<b>Statewide</b>	<b>0.85 (116)</b>	<b>0.91 (115)</b>	<b>+7%</b>	<b>23</b>	<b>33</b>	<b>59</b>

**Table 2.** Drummer densities on the DNR research census areas, 2007-2008.

Area	No. of Drummers (No./100A)	
	2007	2008
Sandhill Hunted (2,020 Acres)	23 (1.1)	24 (1.2)
Sandhill Unhunted (1,300 Acres)	17 (1.3)	19 (1.5)
Stone Lake Exp. Area (3,310 Acres)	71 (2.1)	74 (2.2)



**Figure 1.** Ruffed grouse drumming regions with transect starting points.



**Figure 2.** Mean number of drums/stop on ruffed grouse drumming routes, 1964-2008.